

The impact of different tumour subtypes on management and survival of New Zealand women with Stage I–III breast cancer

Ross Lawrenson, Chunhuan Lao, Ian Campbell, Vernon Harvey, Sanjeewa Seneviratne, Mark Elwood, Diana Sarfati, Marion Kuper-Hommel

ABSTRACT

AIMS: This study aims to describe the prevalence and characteristics of the different ER/PR/HER2 subtypes in New Zealand women with breast cancer, and to explore their treatment and outcomes.

METHODS: This study included women diagnosed with Stage I–III breast cancer between January 2006 and May 2013, recorded in the combined Waikato and Auckland Breast Cancer Registers, and with complete data on their ER, PR and HER2 status. Five ER/PR/HER2 phenotypes were classified. Kaplan-Meier method and Cox proportional hazards model were used to examine the survival differences among these subtypes.

RESULTS: Of the 6,875 eligible women, 4,274 (62.2%) were classified as Luminal A, 836 (12.2%) as Luminal B HER2-, 605 (8.8%) as Luminal B HER2+, 401 (5.8%) as HER2+ non-Luminal and 759 (11.0%) as Triple Negative. Māori and Pacific women were less likely to have Triple Negative disease, while Pacific women were more likely to be HER2+ non-Luminal. The five-year breast cancer-specific survival was worst for HER2+ non-Luminal (80.1%) and Triple Negative (81.9%), followed by Luminal B HER2- (89.3%) and Luminal B HER2+ (91.6%), and was the best for Luminal A (96.8%). The adjusted breast cancer-specific mortality hazard ratio for Triple Negative and HER2+ non-Luminal compared to Luminal A was 4.91 (95% CI: 3.86–6.26) and 3.94 (95% CI: 2.94–5.30), respectively.

CONCLUSIONS: The pattern of phenotype in women with Stage I–III breast cancer is similar to the overseas cohorts. Most New Zealand women with Luminal A breast cancer have a very good prognosis, but the less common subtypes have relatively poor outcomes.

Breast cancer outcomes have been shown to be strongly linked not only to patient characteristics, and the extent of disease at diagnosis but also to the presence or absence of hormonal biomarkers.^{1–4} In the 1970s the discovery of an estrogen receptor (ER) led to the finding that only those tumours that were ER positive were sensitive to hormonal treatment.⁵ This led to the routine measurement of ER status and targeting of treatment, and subsequently, the introduction of the measurement of a progesterone receptor (PR). ER and PR receptor status have implications for prognosis—women with breast cancers that are both ER

and PR positive (+) have a better prognosis. ER status in particular, and to a lesser extent PR status currently have a major influence on the choice of systemic treatment.⁶

In New Zealand, ER and PR status have been routinely measured for the last 25 years. The measurement of human epidermal growth factor receptor 2 (HER2) status became increasingly common from the first part of this century and has been routine since 2006. This is in line with many other countries.^{7,8} In 2006, 12 months of adjuvant therapy was licensed by the FDA for the treatment of Stage I–III HER2 positive breast cancer.⁹ PHARMAC approved funding

a nine-week course of trastuzumab for Stage I–III breast cancer from July 2007,¹⁰ and a 12-month course from July 2010.^{11,12} Other biomarkers such as Ki67, or BRCA gene mutation status, are not routinely measured at this time but may become more relevant in the future.^{13,14} It has become common practice to categorise cancer into conceptual molecular classes that have different prognostic features, and predict response to specific therapies. This has led to a more personalised approach to treatment based on a patient's molecular phenotype. While a number of studies have been published on the prevalence of individual biomarkers in different ethnic groups in New Zealand,^{1,15–17} there has been little opportunity to look at different molecular categories and how they influence treatment or patient outcomes.

The aim of this study was to describe the prevalence and characteristics of the different breast cancer tumour types as indicated by these biomarkers in women with Stage I–III breast cancer. We then looked at the treatment of these women including the use of endocrine therapy, chemotherapy and trastuzumab for breast cancers that were HER2+. Finally we wanted to examine the outcomes in these different groups of breast cancers.

Methods

The studied population have been identified from the combined Waikato and Auckland Breast Cancer Registers.¹⁸ It has clinical details of 12,372 women diagnosed with invasive breast cancer between June 2000 and May 2013. Only women who were diagnosed with Stage I–III breast cancer between January 2006 and May 2013 and had complete data on their ER, PR and HER2 status were included in this study, as HER2 status testing has been routine since 2006.

The register's data includes: 1) patient characteristics: age and ethnicity; 2) tumour information: diagnosis date, cancer stage and biomarkers, and 3) information on treatment: surgery, chemotherapy, trastuzumab, endocrine therapy and radiation therapy. Information on comorbidities has been obtained by reviewing linked data from the National Minimum dataset (NMDS) and characterising patients using the C3 comorbidity index: 1) less or equal to zero, 2) greater than zero but less or equal to one, and 3) greater than one.^{19,20}

In this study, HER2+ was defined as FISH amplified or IHC 3+ according to the 2013 American Society of Clinical Oncology (ASCO) guideline.²¹ Recommended in the 2001 St. Gallen Consensus, ER+ or PR+ was assessed as IHC positive (1+).²² Based on whether the three biomarkers ER, PR and HER2 were either positive or negative, there were eight possible groups defined by ER, PR and HER2 status. We reduced these groups to five categories based on the St. Gallen Consensus recommendation^{3,23,24} and clinical advice and practice in our region. The most common finding in women with breast cancer is a cancer that is both ER and PR positive but HER2 negative. These breast cancers were categorised as Luminal A. Luminal B HER2- includes women whose breast cancer is ER+, but PR- and HER2-. This group is important as women with breast cancers that are PR negative have a poorer prognosis. There is also a small group (1%) of women with breast cancer that is ER- but PR+. We have included these cases in Luminal B HER2-. A further category is women with breast cancers that are HER2+. These women are usually offered adjuvant chemotherapy plus trastuzumab. These women can be divided into those who would benefit from endocrine therapy (ie, breast cancers that are ER+ or ER- but PR+ (Luminal B HER2+) and a second group of breast cancers that are ER-, PR-, but HER2+ (HER2+ non-Luminal)). Finally there is a group that are Triple Negative (ER-, PR-, and HER2-).

Patient outcomes include breast cancer-specific survival and all-cause survival. These mortality data were derived from the New Zealand National Mortality Collection and linked by the National Health Index (NHI) number to the register data. The NHI number is a unique identifier for people who use health and disability services in New Zealand. For all-cause survival analyses, patients without mortality information were considered to be censored on the last updated date for Mortality Collection which was 31 December 2014. For cancer-specific analyses, deaths from other causes were censored on the date of death. Kaplan-Meier method was used to examine the breast cancer-specific survival in the five subtypes. We used Cox proportional hazards model to estimate the hazard ratio of breast cancer-specific mortality and

Table 1: Demographics of patients by cancer subtype.

Sub-groups	Luminal A		Luminal B HER2-		Luminal B HER2+		HER2+ non-Luminal		Triple Negative		P-value for Chi-square test	Total
Region											<0.001	
Auckland	3,421	63.4%	563	10.4%	424	7.9%	342	6.3%	647	12.0%		5,397
Waikato	853	57.7%	273	18.5%	181	12.2%	59	4.0%	112	7.6%		1,478
Year of diagnosis											<0.001	
2006–2007	963	58.0%	221	13.3%	142	8.6%	117	7.1%	216	13.0%		1,659
2008–2009	1,167	62.4%	209	11.2%	156	8.3%	106	5.7%	232	12.4%		1,870
2010–2011	1,202	63.9%	226	12.0%	161	8.6%	97	5.2%	194	10.3%		1,880
2012–2013	942	64.3%	180	12.3%	146	10.0%	81	5.5%	117	8.0%		1,466
Age (years)											<0.001	
<40	179	43.3%	39	9.4%	67	16.2%	45	10.9%	83	20.1%		413
40–49	1,044	65.0%	125	7.8%	169	10.5%	107	6.7%	161	10.0%		1,606
50–59	1,135	61.5%	209	11.3%	179	9.7%	130	7.0%	192	10.4%		1,845
60–69	1,102	63.8%	252	14.6%	125	7.2%	80	4.6%	167	9.7%		1,726
70–79	510	62.9%	124	15.3%	44	5.4%	29	3.6%	104	12.8%		811
80+	304	64.1%	87	18.4%	21	4.4%	10	2.1%	52	11.0%		474
Ethnicity											<0.001	
Others	3,539	61.8%	730	12.7%	479	8.4%	306	5.3%	672	11.7%		5,726
Māori	454	64.2%	83	11.7%	75	10.6%	40	5.7%	55	7.8%		707
Pacific	281	63.6%	23	5.2%	51	11.5%	55	12.4%	32	7.2%		442
Total	4,274	62.2%	836	12.2%	605	8.8%	401	5.8%	759	11.0%		6,875

all-cause mortality by subtype, ER status, PR status, HER2 status and lymph nodes after adjustment for age, ethnicity, stage, comorbidity and year of diagnosis. All data analyses were performed in IBM SPSS statistics 23 (New York, US).

Results

Of the 12,372 invasive breast cancer cases, 574 were metastatic at diagnosis and 11,798 were Stage I–III at diagnosis. Of the Stage I–III breast cancer cases, 4,475 cases were diagnosed in 2000–2005 and 7,320 were diagnosed in 2006–2013. Of those 7,320 cases diagnosed in 2006–2013, 448 (6.1%) without complete ER, PR or HER2 results were excluded from this study. Those 6,875 women who had complete information on their ER, PR and HER2 status were included.

Of the included cancer cases, 4,274 (62.2%) cases were classified as Luminal A, 836 (12.2%) as Luminal B HER2-, 605

(8.8%) as Luminal B HER2+, 401 (5.8%) as HER2+ non-Luminal and 759 (11.0%) as Triple Negative (Table 1). The mean age varied by subgroup from 58.5 years in Luminal A, 61.1 years in Luminal B HER2-, 54.1 years in Luminal B HER2+, 53.5 years in HER2+ non-Luminal and 57.2 years in Triple Negative. Women with breast cancers that were HER2+ or Triple Negative breast cancer were younger than those classified as Luminal A or Luminal B HER2-. Māori and Pacific women were more likely to have HER2+ breast cancer but less likely to have Triple Negative disease than non-Māori/non-Pacific women. There were stark differences in stage and grade of cancer at diagnosis between the different subtypes: 32.4% of women in HER2+ non-Luminal had Stage III cancer compared to 12.5% in Luminal A; 80.8% of women with Triple Negative cancer had Grade 3 disease while only 12.0% in Luminal A had Grade 3 cancer (Table 2).

Table 2: Tumour characteristics and treatment by cancer subtype.


Subgroups	Luminal A		Luminal B HER2-		Luminal B HER2+		HER2+ non-Luminal		Triple Negative		P-value for Chi-square test	Total	
Tumour size (mm)											<0.001		
0~10	720	17.1%	133	16.5%	70	11.9%	71	18.3%	84	11.2%		1,078	16.0%
10~20	1,661	39.5%	251	31.1%	187	31.7%	93	24.0%	220	29.3%		2,412	35.8%
20~30	931	22.2%	208	25.7%	158	26.8%	82	21.2%	217	28.9%		1,596	23.7%
30~50	603	14.4%	154	19.1%	115	19.5%	86	22.2%	178	23.7%		1,136	16.9%
50+	286	6.8%	62	7.7%	60	10.2%	55	14.2%	53	7.0%		516	7.7%
Unknown	73		28		15		14		7			137	
Stage											<0.001		
I	2,223	52.0%	358	42.8%	214	35.4%	121	30.2%	268	35.3%		3,184	46.3%
II	1,515	35.4%	346	41.4%	245	40.5%	150	37.4%	358	47.2%		2,614	38.0%
III	536	12.5%	132	15.8%	146	24.1%	130	32.4%	133	17.5%		1,077	15.7%
Grade											<0.001		
1	1,545	36.7%	190	23.3%	30	5.1%	5	1.3%	17	2.3%		1,787	26.5%
2	2,164	51.4%	374	45.9%	282	47.5%	74	19.1%	126	16.9%		3,020	44.7%
3	504	12.0%	250	30.7%	282	47.5%	309	79.6%	603	80.8%		1,948	28.8%
Unknown	61		22		11		13		13			120	
Lymph nodes											<0.001		
No positive lymph nodes	2,601	63.8%	479	60.6%	296	50.7%	175	45.3%	462	62.4%		4,013	61.0%
Positive lymph nodes	1,479	36.3%	312	39.4%	288	49.3%	211	54.7%	278	37.6%		2,568	39.0%
Unknown	194		45		21		15		19			294	
Surgery											<0.001		
Breast conserving surgery	2,520	59.0%	424	50.7%	267	44.1%	132	32.9%	361	47.6%		3,704	53.9%
Mastectomy	1,650	38.6%	384	45.9%	324	53.6%	259	64.6%	390	51.4%		3,007	43.7%
No primary surgery	104	2.4%	28	3.3%	14	2.3%	10	2.5%	8	1.1%		164	2.4%
Endocrine therapy											<0.001		
No endocrine therapy	1210	28.3%	213	25.5%	75	12.4%	381	95.0%	723	95.3%		2,602	37.8%
Endocrine therapy	3,064	71.7%	623	74.5%	530	87.6%	20	5.0%	36	4.7%		4,273	62.2%
Chemotherapy											<0.001		
No chemotherapy	3,302	77.3%	575	68.8%	173	28.6%	92	22.9%	252	33.2%		4,394	63.9%
Chemotherapy	972	22.7%	261	31.2%	432	71.4%	309	77.1%	507	66.8%		2,481	36.1%
Trastuzumab											<0.001		
No trastuzumab	4,264	99.8%	832	99.5%	204	33.7%	106	26.4%	750	98.8%		6,156	89.5%
Trastuzumab	10	0.2%	4	0.5%	401	66.3%	295	73.6%	9	1.2%		719	10.5%
Total	4,274		836		605		401		759			6,875	

Table 3: Five-year breast cancer-specific survival and all-cause survival by subtype.

Subtype	Breast cancer-specific survival		All-cause survival	
	5-year survival (95% CI)		5-year survival (95% CI)	
Luminal A	96.8%	(96.2%–97.4%)	91.9%	(90.9%–92.8%)
Luminal B HER2-	89.3%	(86.9%–91.8%)	81.6%	(78.6%–84.7%)
Luminal B HER2+	91.6%	(89.0%–94.2%)	87.5%	(84.4%–90.6%)
HER2+ non-Luminal	80.1%	(75.6%–84.6%)	78.1%	(73.5%–82.7%)
Triple Negative	81.9%	(78.9%–84.9%)	76.7%	(73.4%–79.9%)

CI: confidence interval

Figure 1: Breast cancer-specific survival by subtype using the Kaplan Meier method.



Follow-up time (months)		0	12	24	36	48	60	72	84	96
Number of women at risk	Luminal A	4,274	4,241	3,916	3,209	2,578	1,987	1,413	866	452
	Luminal B HER2-	836	814	733	582	460	349	261	170	88
	Luminal B HER2+	605	596	549	436	355	267	192	125	62
	HER2+ non-Luminal	401	394	350	277	217	169	125	85	40
	Triple Negative	759	738	651	544	420	342	260	157	80

As expected, the treatment varied depending on the subtype identified. In total, 97.6% of women were treated with surgery, and women with Luminal A were more likely to be treated with breast conserving surgery. In contrast, 64.6% of women with ER-, PR- and HER2+ breast cancer (HER2+ non-Luminal) were treated with a mastectomy. Of Luminal A women, 71.7% received endocrine therapy compared to 87.6% of the women with Luminal B HER2+ cancer. Chemotherapy was more likely to be prescribed for breast cancers with the worst prognosis, ie, cancers that were HER2+ or Triple Negative. Of the cancers that were HER2+, those who were ER- and PR- (HER2+ non-Luminal) were more likely to receive trastuzumab than those who were ER/PR positive (Luminal B HER2+).

Overall Luminal A women had a very good prognosis while women with cancers that were HER2+, ER- or were Triple Negative had a relatively poor prognosis (Figure 1). The five-year breast cancer-specific survival (Table 3) was worst for HER2+ non-Luminal

(80.1%) and Triple Negative (81.9%), followed by Luminal B HER2- (89.3%) and Luminal B HER2+ (91.6%), and was the best for Luminal A (96.8%, Log-rank test p-value <0.001).

After adjustment for age, ethnicity, stage, comorbidity and year of diagnosis, women with Triple Negative breast cancer had the worst prognosis (Table 4): hazard ratio of 4.91 (95% CI: 3.86–6.26, p-value<0.001) for breast cancer-specific mortality and 2.74 (95% CI: 2.29–3.28, p-value<0.001) for all-cause mortality compared to Luminal A. The second worst prognosis was HER2+ non-Luminal with a hazard ratio of 3.94 (95% CI:2.94–5.30, p-value<0.001) for breast cancer-specific mortality and 2.46 (95% CI:1.92–3.15, p-value<0.001) for all-cause mortality compared to Luminal A. Breast cancer-specific mortality hazard ratios were 3.19 (95% CI: 2.65–3.85, p-value<0.001) for ER-, 3.29 (95% CI:2.72–3.98, p-value<0.001) for PR-, 1.58 (95% CI:1.28–1.96, p-value<0.001) for HER2+ and 1.18 (95% CI:0.89–1.55, p-value=0.248) for lymph node positive, respectively.

Table 4: Hazard ratios in breast cancer-specific mortality and all-cause mortality after adjustment for age, ethnicity, stage, comorbidity and year of diagnosis.

Subgroups	Breast cancer-specific mortality			All-cause mortality		
	Hazard ratio	95% CI	p-value	Hazard ratio	95% CI	p-value
Subtype						
Luminal A	Ref			Ref		
Luminal B HER2-	2.64	1.98–3.51	<0.001	1.85	1.52–2.25	<0.001
Luminal B HER2+	2.04	1.47–2.82	<0.001	1.46	1.14–1.88	0.003
HER2+ non-Luminal	3.94	2.94–5.30	<0.001	2.46	1.92–3.15	<0.001
Triple Negative	4.91	3.86–6.26	<0.001	2.74	2.29–3.28	<0.001
ER status						
ER+	Ref			Ref		
ER-	3.19	2.65–3.85	<0.001	2.21	1.91–2.56	<0.001
PR status						
PR+	Ref			Ref		
PR-	3.29	2.72–3.98	<0.001	2.12	1.85–2.43	<0.001
HER2 status						
HER2-	Ref			Ref		
HER2+	1.58	1.28–1.96	<0.001	1.36	1.14–1.62	<0.001
Lymph nodes						
No positive lymph nodes	Ref			Ref		
Positive lymph nodes	1.18	0.89–1.55	0.248	1.08	0.88–1.32	0.480

Discussion

The proportion of women in each subgroup was similar to that found in other large studies with 62% women having Luminal A tumours and 11% having Triple Negative type tumours.²⁵ The differences in characteristics by subtype in our study are also consistent with other international cohorts.^{2,26} As age increased, the proportion of women with Luminal A breast cancer increased. HER2+ non-Luminal and Triple Negative breast cancers were more likely to be Grade 3 (80%), and Luminal A cancers were the least likely to be Grade 3 (12%). HER2+ cancers were more likely to have positive lymph nodes and worse cancer stage than other subtype cancers.^{2,26}

The findings from two large cancer centres in New Zealand show that treatment for Stage I–III breast cancer is tailored to the subtype with variation in the use of endocrine therapy, chemotherapy and trastuzumab. Women with rarer subtypes such as HER2+ and Triple Negative were more likely to receive chemotherapy and when identified either endocrine therapy or trastuzumab. On the other hand, women with Luminal A disease who have a good prognosis were less likely to receive chemotherapy. Surgical treatment also varied by subtype. Women with Luminal A cancer were more likely to be treated with breast conserving surgery. However, women with phenotypes with poor prognosis were less likely to receive breast conserving surgery. No doubt this is affected by the prognosis of the subtype, but other factors such as the size of the tumour, lymph node involvement, stage and grade also affect surgical treatment.²⁷

As well as noting the different characteristics and treatment of women at the time of diagnosis in the five subgroups there were also differences in outcomes. The survival curves show that in the majority of women, ie, those in Luminal A, the five-year survival was 97%, while for those with ER and PR negative, HER2+ disease only 80% survive five years. Having a cancer that was either ER or PR negative was also an important prognostic indicator, with a hazard ratio for ER negative of 3.19 and for PR negative of 3.29. Women with HER2+ or Triple Negative disease and are more likely to be younger and have Grade 3 disease. On the other hand, women with Luminal A disease are likely to be older and do better.

This is consistent with the literature.^{2,28} We also know that in New Zealand outcomes for Māori and Pacific women are poor.^{1,29–31} While they may be slightly more likely to have HER2 positive disease, they are less likely to have the subtype with the worst prognosis, ie, Triple Negative disease. It has been shown that for Māori the differences in biology only make a small contribution to the differences in outcomes.¹

The strength of this study is that it comprises a relatively large population-based database with comprehensive data on patient characteristics, patient treatment as well as outcomes. One weakness is that we did not take into account other important biomarkers such as Ki67. We also have not included grade of disease in our classification. Some classification systems would classify ER+, PR+ and HER2- breast cancers as luminal B rather than luminal A if they are high grade or have a high Ki67,^{32–34} but these were all classified into Luminal A in our study. Doing the classification this way would slightly bias luminal A cases towards worse outcomes in our study. Our classification may differ from a classification based on gene expression profiling. On the other hand, our grouping of cancers into five subtypes is also a strength of this study as breast cancer treatment decisions are generally based on the presence or absence of these three biomarkers. Gene expression profiling is not routinely available in clinical practice and only infrequently used to assist treatment decisions in New Zealand at present.

Conclusions

The pattern of phenotype in women with Stage I–III breast cancer is similar to the international cohorts. Most New Zealand women with Luminal A breast cancer have a very good prognosis, but the less common subtypes have relatively poor outcomes. We have demonstrated differences in tumour grade, stage, patient age and ethnicity according to breast cancer subtype in a New Zealand population. The treatment of women with Stage I–III breast cancer varies by molecular phenotype. Treatment is becoming personalised to their individual molecular phenotype. Despite this there was a major variation in the prognosis of women with Stage I–III breast cancer with differing molecular phenotype.

Competing interests:

RL, SS, IC, ME, CL and DS report grants from Health Research Council of New Zealand during the conduct of the study.

Acknowledgements:

We would like to acknowledge the financial support from the Health Research Council of New Zealand, the Auckland and Waikato Breast Cancer Registers for providing the detailed data, and the New Zealand Breast Cancer Foundation and the Waikato Bay of Plenty Division of the Cancer Society for funding the Registers.

This study was supported by the Health Research Council of New Zealand for the project 'Improving outcomes for women with breast cancer in NZ' (grant number: 14/484).

The funding source did not have any role in the design of the study; in the analysis and interpretation of the data; or in the preparation of the manuscript.

Author information:

Ross Lawrenson, Professor of Population Health, Waikato Medical Research Centre, The University of Waikato, Hamilton; Chunhuan Lao, Research Fellow, Waikato Medical Research Centre, The University of Waikato, Hamilton; Ian Campbell, Associate Professor of Surgery, School of Medicine, The University of Auckland, Auckland; Vernon Harvey, Medical Oncologist, Auckland City Hospital, Auckland; Sanjeewa Seneviratne, Senior Lecturer, Department of Surgery, Faculty of Medicine, University of Colombo, Sri Lanka; Mark Elwood, Professor of Cancer Epidemiology, School of Population Health, The University of Auckland, Auckland; Diana Sarfati, Professor of Epidemiology, Department of Public Health, The University of Otago, Wellington; Marion Kuper-Hommel, Medical Oncologist, Waikato District Health Board, Hamilton.

Corresponding author:

Prof Ross Lawrenson, University of Waikato, Level 3 Hockin Building, Waikato Hospital, Hamilton 3240.

ross.lawrenson@waikatodhb.health.nz

URL:

<http://www.nzma.org.nz/journal/read-the-journal/all-issues/2010-2019/2018/vol-131-no-1475-18-may-2018/7572>

REFERENCES:

1. Seneviratne S, Campbell I, Scott N, Shirley R, Peni T, Lawrenson R. Ethnic differences in breast cancer survival in New Zealand: contributions of differences in screening, treatment, tumor biology, demographics and comorbidities. *Cancer Causes and Control* 2015; 26:1813–24.
2. Parise CA, Caggiano V. Breast cancer survival defined by the er/pr/her2 subtypes and a surrogate classification according to tumor grade and immunohistochemical biomarkers. *Journal of Cancer Epidemiology* 2014 ;Vol 2014;469251.
3. Goldhirsch A, Winer EP, Coates AS, et al. Personalizing the treatment of women with early breast cancer: Highlights of the st gallen international expert consensus on the primary therapy of early breast Cancer 2013. *Annals of Oncology* 2013; 24:2206–23.
4. Early Breast Cancer Trialists' Collaborative Group. Relevance of breast cancer hormone receptors and other factors to the efficacy of adjuvant tamoxifen: Patient-level meta-analysis of randomised trials. *The Lancet* 2011; 378:771–84.
5. Block GE, Jensen EV, Polley Jr TZ. The prediction of hormonal dependency of mammary cancer. *Annals of Surgery* 1975; 182:342–52.
6. Seneviratne S, Campbell I, Scott N, et al. Adherence to adjuvant endocrine therapy: Is it a factor for ethnic differences in breast cancer outcomes in New Zealand? *Breast* 2015; 24:62–7.
7. Duffy MJ, O'Donovan N, McDermott E, Crown J. Validated biomarkers: The key to precision treatment in patients with breast cancer. *Breast* 2016; 29:192–201.
8. Anderson WF, Pfeiffer RM, Wohlfahrt J, Ejlersen B, Jensen MB, Kroman N. Associations of pari-

- ty-related reproductive histories with ER+/- and HER2+/- receptor-specific breast cancer aetiology. *International journal of epidemiology* 2017;46:373.
9. FDA Approval for Trastuzumab. 2013. at <http://www.cancer.gov/about-cancer/treatment/drugs/fda-trastuzumab>
 10. Metcalfe S, Evans J, Priest G. PHARMAC funding of 9-week concurrent trastuzumab (Herceptin) for HER2-positive early breast cancer. *New Zealand Medical Journal* 2007; 120(1256).
 11. Govt dodges Pharmac to fund full Herceptin courses. 2008. (Accessed 11 July, 2016)
 12. Lawrenson R, Lao C, Campbell I, et al. The use of trastuzumab in New Zealand women with breast cancer. *Asia-Pacific Journal of Clinical Oncology* 2018; Apr 14(2):e152–e160.
 13. Yerushalmi R, Woods R, Ravdin PM, Hayes MM, Gelmon KA. Ki67 in breast cancer: prognostic and predictive potential. *The Lancet Oncology* 2010; 11:174–83.
 14. Farmer H, McCabe H, Lord CJ, et al. Targeting the DNA repair defect in BRCA mutant cells as a therapeutic strategy. *Nature* 2005; 434:917–21.
 15. Seneviratne S, Lawrenson R, Scott N, Kim B, Shirley R, Campbell I. Breast cancer biology and ethnic disparities in breast cancer mortality in New Zealand: A cohort study. *PLoS ONE* 2015; 10(4), e0123523.
 16. Seneviratne S, Lawrenson R, Harvey V, et al. Stage of breast cancer at diagnosis in New Zealand: Impacts of socio-demographic factors, breast cancer screening and biology. *BMC Cancer* 2016; 16(1):129.
 17. Davey V, Robinson B, Dijkstra B, Harris G. The Christchurch Breast Cancer Patient Register: The first year. *New Zealand Medical Journal* 2012; 125:37–47.
 18. Seneviratne S, Campbell I, Scott N, Shirley R, Peni T, Lawrenson R. Accuracy and completeness of the New Zealand Cancer Registry for staging of invasive breast cancer. *Cancer Epidemiology* 2014; 38:638–44.
 19. Tin ST, Elwood JM, Lawrenson R, Campbell I, Harvey V, Seneviratne S. Differences in breast cancer survival between public and private care in New Zealand: Which factors contribute? *PLoS ONE* 2016; 11 (4), e1053206.
 20. Sarfati D, Gurney J, Stanley J, et al. Cancer-specific administrative data-based comorbidity indices provided valid alternative to Charlson and National Cancer Institute Indices. *Journal of Clinical Epidemiology* 2014; 67:586–95.
 21. Wolff AC, Hammond ME, Hicks DG, et al. Recommendations for human epidermal growth factor receptor 2 testing in breast cancer: American Society of Clinical Oncology/College of American Pathologists clinical practice guideline update. *J Clin Oncol* 2013; 31:3997–4013.
 22. Thuerlimann B. International consensus meeting on the treatment of primary breast cancer 2001, St. Gallen, Switzerland. *Breast cancer (Tokyo, Japan)* 2001; 8:294–7.
 23. Coates AS, Winer EP, Goldhirsch A, Gelber RD, Gnant M, Piccart-Gebhart M. Panel Members. Tailoring therapies-improving the management of early breast cancer: St Gallen International Expert Consensus on the Primary Therapy of Early Breast Cancer 2015. *Ann Oncol* 2015; 26(8):1533–1546.
 24. Goldhirsch A, Wood WC, Coates AS, Gelber RD, Thürlimann B, Senn HJ. Strategies for subtypes-dealing with the diversity of breast cancer: Highlights of the St Gallen international expert consensus on the primary therapy of early breast cancer 2011. *Annals of Oncology* 2011; 22:1736–47.
 25. Parise C, Caggiano V. Breast cancer mortality among Asian-American women in California: Variation according to ethnicity and tumor subtype. *Journal of Breast Cancer* 2016; 19:112–21.
 26. Voduc KD, Cheang MC, Tyldesley S, Gelmon K, Nielsen TO, Kennecke H. Breast cancer subtypes and the risk of local and regional relapse. *J Clin Oncol* 2010; 28:1684–91.
 27. Seneviratne S, Scott N, Lawrenson R, Campbell I. Ethnic, socio-demographic and socio-economic differences in surgical treatment of breast cancer in New Zealand. *ANZ Journal of Surgery* 2017; 87(7–8):E32–E39.
 28. O'Brien KM, Cole SR, Tse CK, et al. Intrinsic breast tumor subtypes, race, and long-term survival in the carolina breast cancer study. *Clinical Cancer Research* 2010; 16:6100–10.
 29. Bennett H, Marshall R, Campbell I, Lawrenson R. Women with breast cancer in Aotearoa New Zealand: The effect of urban versus rural residence on stage at diagnosis and survival. *New Zealand Medical Journal* 2007; 120(1266).
 30. Lawrenson R, Lao C, Campbell I, et al.

- Treatment and survival disparities by ethnicity in New Zealand women with stage I-III breast cancer tumour subtypes. *Cancer causes & control* : CCC 2017; 28:1417–27.
31. Lawrenson R, Seneviratne S, Scott N, Peni T, Brown C, Campbell I. Breast cancer inequities between Maori and non-Maori women in Aotearoa/New Zealand. *European journal of cancer care* 2016; 25:225–30.
32. Metzger-Filho O, Sun Z, Viale G, et al. Patterns of recurrence and outcome according to breast cancer subtypes in lymph node-negative disease: Results from international breast cancer study group trials VIII and IX. *Journal of Clinical Oncology* 2013; 31:3083–90.
33. Laurberg T, Alsner J, Tramm T, et al. Impact of age, intrinsic subtype and local treatment on long-term local-regional recurrence and breast cancer mortality among low-risk breast cancer patients. *Acta Oncologica* 2017; 56:59–67.
34. Von Minckwitz G, Untch M, Blohmer JU, et al. Definition and impact of pathologic complete response on prognosis after neoadjuvant chemotherapy in various intrinsic breast cancer subtypes. *Journal of Clinical Oncology* 2012; 30:1796–804.

Copyright of New Zealand Medical Journal is the property of New Zealand Medical Journal and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.